Children's

Relief you can trust



Find the right dose on the chart below. If possible, use weight

to dose

otherwise

use age.

Infants' Advil **Concentrated Drops** for ages 6-23 months

ACTIVE INGREDIENT

Fever reducer; relieves minor aches and pains due to the common cold, flu, headaches, and toothaches



Children's Advil Suspension for ages 2-11 years

ACTIVE INGREDIENT Ibuprofen 100 mg (NSAID)† in each 5 mL

Fever reducer; relieves minor aches and pains due to the common cold, flu, sore throat headaches, and toothaches



Junior Strength Advil Chewables for ages 6-11 years

ACTIVE INGREDIENT

Fever reducer; relieves minor aches and pains due to the common cold, flu, sore throat headaches, and toothaches



Junior Strength Advil for ages 6-11 years

ACTIVE INGREDIENT Ibuprofen 100 mg (NSAID)† in each tablet

Fever reducer; relieves minor aches and pains due to the common cold, flu, sore throat headaches, and toothaches

	Weight	Age	GIVE YOUR CHILD TH	THE DOSE INDICATED BELOW EVERY 6-8 HOURS, NOT TO EXCEED 4 DOSES/24 HOURS			
PATIENT DOSING	12-17 pounds	under 6 months	Ask a doctor	Ask a doctor about using Infants' Advil			
		6-11 months	1.25 mL‡		Ask a doctor about using Infants' or Children's Advil		
	18-23 pounds	12-23 months	1.875 mL‡				
	24-35 pounds	2-3 years	Ask a doctor about using Children's or Junior Strength Advil	1 tsp [‡]			
	36-47 pounds	4-5 years		1½ tsp‡			
	48-59 pounds	6-8 years		2 tsp‡	2 tablets	2 tablets	
	60-71 pounds	9-10 years		2½ tsp‡	2½ tablets	2 tablets	
	72-95 pounds	11 years		3 tsp‡	3 tablets	3 tablets	
	- 40.		atropath Advil				

For 12+ years, use adult strength Advil

Use the fever diary below to record your child's fevers and dosing schedule.

Fever:	°F or°C	Time:	Dose:
Fever:	°F or°C	Time:	Dose:
Fever:	°F or°C	Time:	Dose:
Fever:	°F or°C	Time:	Dose:

Please refer to Drug Facts on the product label or visit Advil.com for complete warnings. **Important Instructions for Proper Use**

- Read and keep the cartons for complete warnings and information on Advil products
- If you are taking other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), then consult your doctor before taking any Advil products
 Adult strength medicines should not be administered to children
- Measure dosing only with the device that comes with the product



Shake liquid products well before dosing. Visit ChildrensAdvil.com for Drug Facts, product updates and additional information.

[‡]Measure dosing only with the device provided.

[†]Nonsteroidal anti-inflammatory drug.